

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 April 2001 (05.04.2001)

PCT

(10) International Publication Number
WO 01/23588 A1

- (51) International Patent Classification⁷: C12N 15/57, 9/48
- (21) International Application Number: PCT/US00/26876
- (22) International Filing Date:
29 September 2000 (29.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/156,685 29 September 1999 (29.09.1999) US
- (71) Applicant: LEXICON GENETICS INCORPORATED
[US/US]; 4000 Research Forest Drive, The Woodlands, TX
77381 (US).
- (72) Inventors: DONOHO, Gregory; 41 Fallenstone Drive,
The Woodlands, TX 77381 (US). TURNER, C., Alexan-
der, Jr.; 67 Winter Wheat Place, The Woodlands, TX
77381 (US). NEHLS, Michael; Paul-Keller-Strasse 6,
82131 Stockdorf (DE). FRIEDRICH, Glenn; Breland
& Breland, 2207 Hermann Drive, Houston, TX 77004
(US). ZAMBROWICZ, Brian; 18 Firethorne Place, The
Woodlands, TX 77382 (US). SANDS, Arthur, T.; 163
Bristol Bend Circle, The Woodlands, TX 77382 (US).
- (74) Agent: ISHIMOTO, Lance; Lexicon Genetics Incorpo-
rated, 4000 Research Forest Drive, The Woodlands, TX
77381 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— *With international search report.*
— *Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.*
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: HUMAN CARBOXYPEPTIDASES AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that can be used in therapeutic, diagnostic, and pharmacogenomic applications.

WO 01/23588 A1

**NOVEL HUMAN CARBOXYPEPTIDASES AND
POLYNUCLEOTIDES ENCODING THE SAME**

The present application claims the benefit of U.S.
5 Provisional Application Number 60/156,685 which was filed on
September 29, 1999 and is herein incorporated by reference in its
entirety.

1. INTRODUCTION

The present invention relates to the discovery,
10 identification, and characterization of novel human
polynucleotides encoding proteins that share sequence similarity
with animal proteases. The invention encompasses the described
polynucleotides, host cell expression systems, the encoded
proteins, fusion proteins, polypeptides and peptides, antibodies
15 to the encoded proteins and peptides, and genetically engineered
animals that either lack or over express the disclosed genes,
antagonists and agonists of the proteins, and other compounds that
modulate the expression or activity of the proteins encoded by the
disclosed sequences that can be used for diagnosis, drug
20 screening, clinical trial monitoring and the treatment of
physiological disorders.

2. BACKGROUND OF THE INVENTION

Proteases are enzymes that cleave polypeptide sequences.
Carboxypeptidases are proteases that hydrolyze the peptide bonds
25 at the carboxy-terminal end of a chain of amino acids and have
been identified in a wide variety of cell types and animals.
Peptidases have been implicated in a wide variety of biological
processes including, but not limited to, digestion, coagulation,
diabetes, prostate cancer, gynecological disorders, neurological
30 disorders, and obesity. Accordingly, peptidases represent
significant targets for regulatory control of a variety of
physiological processes and pathways.

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery, identification, and characterization of nucleotides that encode novel human proteins, and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs) described for the first time herein share structural similarity with animal proteases, and especially carboxypeptidases. As such, the described NHPs represent a new family of protease-related proteins with a range of homologues and orthologs that transcend phyla and a broad range of species.

The novel human nucleic acid sequences described herein, encode proteins/open reading frames (ORFs) of 47, 88, 247, 92, 437, and 350 amino acids in length (see SEQ ID NOS: 2, 4, 6, 8, 10, and 12 respectively).

The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof that compete with native NHP, peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and gene or regulatory sequence replacement constructs) or to enhance the expression of the described NHP sequences (e.g., expression constructs that place the described sequence under the control of a strong promoter system), and transgenic animals that express a NHP transgene, or "knock-outs" (which can be conditional) that do not express a functional NHP.

Further, the present invention also relates to processes of identifying compounds that modulate, i.e., act as agonists or antagonists, of NHP expression and/or NHP product activity that utilize purified preparations of the described NHPs and/or NHP product, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

The Sequence Listing provides the sequences of 6 protease-related ORFs that encode the described NHP amino acid sequences.

5

5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs, described for the first time herein, are novel proteins that are expressed in, *inter alia*, human cell lines, and human brain, pituitary, spinal cord, thymus, spleen, lymph node, bone marrow, trachea, lung, kidney, prostate, testis, thyroid, adrenal gland, stomach, small intestine colon, skeletal muscle, uterus, mammary gland, bladder, and cervix cells. The described sequences were compiled from gene trapped cDNAs and a clone isolated from a human prostate cDNA library (Edge Biosystems, Gaithersburg, MD). The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described genes, including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of the NHPs that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including but not limited to the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including but not limited to soluble proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of an NHP, or one of its domains (e.g., a receptor/ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide;

or (e) therapeutic or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in the Sequence Listing.

5 As discussed above, the present invention includes: (a) the human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the
10 Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current
15 Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent gene product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of the DNA sequence that encode and express an amino acid sequence presented in the Sequence Listing
20 under moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, supra), yet still encode a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or
25 engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No. 5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding NHP
30 ORFs, or their functional equivalents, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85 percent similar to corresponding regions of SEQ ID NO:1 (as measured by BLAST

sequence comparison analysis using, for example, the GCG sequence analysis package using default parameters).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP gene nucleotide sequences. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about 20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length may partially overlap each other and/or the NHP sequence may be represented using oligonucleotides that do not overlap. Accordingly, the described NHP polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 18, and preferably about 25, nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences may begin at any nucleotide present within a sequence in the Sequence Listing and proceed in either a sense (5'-to-3')

orientation vis-a-vis the described sequence or in an antisense orientation.

For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP sequence antisense molecules, useful, for example, in NHP sequence regulation (for and/or as antisense primers in amplification reactions of NHP nucleic acid sequences). With respect to NHP sequence regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple helix sequences that are also useful for NHP sequence regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641).

The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example,

Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual (and periodic updates thereof), Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.

5 Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, 10 nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene 15 can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

 Further, a NHP sequence homolog can be isolated from nucleic 20 acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA 25 prepared from, for example, human or non-human cell lines or tissue known or suspected to express an allele of a NHP sequence.

 The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP 30 gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment

can be used to isolate genomic clones via the screening of a genomic library.

PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard
5 procedures, from an appropriate cellular or tissue source (i.e., one known, or suspected, to express a NHP sequence, such as, for example, testis tissue). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the
10 priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be
15 isolated. For a review of cloning strategies that can be used, see e.g., Sambrook et al., 1989, *supra*.

A cDNA encoding a mutant NHP sequence can be isolated, for example, by using PCR. In this case, the first cDNA strand can be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA
20 isolated from tissue known or suspected to be expressed in an individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene.
25 Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s)
30 responsible for the loss or alteration of function of the mutant NHP sequence product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a

mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, obesity, high blood pressure, etc.), or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant NHP allele. A normal NHP sequence, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP coding sequences can then be purified and subjected to sequence analysis according to methods well known to those skilled in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant NHP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

Additionally, screening can be accomplished using labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP sequence product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known in the art.

The invention also encompasses (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a

regulatory element that directs the expression of the coding sequences (for example, baculo virus as described in U.S. Patent No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP sequence under the control of an exogenously introduced regulatory element (i.e., gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the human cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of the NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP sequence (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs or inappropriately expressed NHPs for

the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs
5 (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of NHP in the body. The use of engineered host cells and/or animals may offer an advantage in that such systems allow not only for the
10 identification of compounds that bind to an endogenous receptor/ligand of a NHP, but can also identify compounds that trigger NHP-mediated activity.

Finally, the NHP products can be used as therapeutics. For example, soluble versions or derivatives of a NHP, or
15 peptides/domains corresponding a NHP, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in
20 a NHP-mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics the NHP could activate or effectively antagonize NHP function. Nucleotide constructs
25 encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding functional
30 NHPs, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses

pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

5

5.1 THE NHP SEQUENCES

The cDNA sequences and the corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing. The NHP sequences were obtained from a human prostate
10 cDNA library using probes and/or primers generated from human gene trapped sequence tags. Expression analysis has provided evidence that the described NHPs can be expressed, for example, in a variety of human cell types and that the described NHPs share
15 carboxypeptidase B or carboxypeptidase A, from, *inter alia*, humans, mice, and rats.

SEQ ID NO: 13 describes a full length NHP ORF with flanking 5' and 3' sequences.

5.2 NHPS AND NHP POLYPEPTIDES

20 NHPs, polypeptides, peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as reagents in diagnostic assays, for the identification of other cellular gene
25 products related to a NHP, as reagents in assays for screening for compounds that can be as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and disease. Several uses and applications for plasma carboxypeptidases similar to those described herein are described
30 in U.S. Patent No. 5,593,674, the disclosure of which is herein incorporated by reference in its entirety.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP polynucleotides. The NHPs have

initiator methionines in DNA sequence contexts consistent with a translation initiation site and a hydrophobic signal-like sequence is present near the N-terminal region of the protein. The sequence data presented herein indicate that alternatively spliced
5 forms of the NHPs exist (which may or may not be tissue specific).

The NHP amino acid sequences of the invention include the nucleotide and amino acid sequences presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by
10 the invention. In fact, any NHP protein encoded by the NHP nucleotide sequences described above, are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is
15 well known, and, accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken
20 together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al. eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can
25 encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a
30 substrate of a NHP, or the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins

include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described above, but which result in a silent change, thus producing a functionally equivalent gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. The presently described NHPs are similar to plasma carboxypeptidase B and are likely soluble proteins. Where the NHP peptide or polypeptide to be expressed is a soluble NHP protein, or a NHP peptide derived from a substantially nonhydrophobic domain of a NHP, or a truncated or deleted NHP the peptide or polypeptide can be recovered from the culture, i.e., from the host cell in cases where the NHP peptide or polypeptide is not secreted, or from the culture media in cases where the NHP peptide or polypeptide is secreted by the cells. However, such expression systems also encompass engineered host cells that express a NHP, or functional equivalent, *in situ*, i.e., anchored in the cell membrane. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural

and functional characteristics of the NHP, but to assess biological activity, e.g., in-drug screening assays.

The expression systems that can be used for purposes of the invention include but are not limited to microorganisms such as
5 bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences; insect cell systems infected
10 with recombinant virus expression vectors (e.g., baculovirus) containing NHP sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid)
15 containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus
20 7.5K promoter).

In bacterial systems, a number of expression vectors can be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of
25 pharmaceutical compositions of or containing NHP, or for raising antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J.
30 2:1791), in which a NHP coding sequence may be ligated individually into the vector in frame with the *lacZ* coding region so that a fusion protein is produced; pIN vectors (Inouye &

Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target sequence product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhydrosis virus (AcNPV) is used as a vector to express foreign sequences. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted sequence is expressed (e.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of

expressing a NHP product in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the

5 ATG initiation codon and adjacent sequences. In cases where an entire NHP sequence or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding

10 sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control

15 signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bittner et al., 1987, Methods in Enzymol. 153:516-544).

20 In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the sequence product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of

25 the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To

30 this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include, but are not limited to,

CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the NHP sequences described above may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk⁻, hgp⁺ or ap⁺ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers

resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hygromycin (Santerre, et al., 1984, Gene 30:147).

Alternatively, any fusion protein may be readily purified by
5 utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88: 8972-8976). In this system, the sequence
10 of interest is subcloned into a vaccinia recombination plasmid such that the sequence's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose columns and
15 histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes
20 of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments,
25 fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Antibodies, and applications are uses thereof, similar to those contemplated herein are described in U.S. Patent No. 5,474,901 the disclosure of which is herein incorporated by
30 reference in its entirety.

The antibodies of the invention can be used, for example, in the detection of NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby

patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes, as described, below, in Section 5.5, for the evaluation of the effect of test compounds on expression and/or activity of a NHP coding sequence product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into the patient. Such antibodies may additionally be used as a method for the inhibition of abnormal NHP activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with the NHP, an NHP peptide (e.g., one corresponding to a functional domain of an NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP or mutated variant of the NHP. Such host animals may include but are not limited to pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger et al., 1984, Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patents Nos. 6,075,181 and 5,877,397 and their respective disclosures which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 334:544-546) can be adapted to produce single chain antibodies against NHP sequence

products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, J. Immunol. 147(8):2429-2438). For example antibodies which bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor/ligand can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor, cofactor, ligand, or binding partner. Such anti-idiotypic antibodies or Fab fragments of such anti-idiotypes can be used in therapeutic regimens involving a NHP mediated pathway.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents,

and patent applications are herein incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 24 contiguous bases of nucleotide sequence first disclosed in the NHP polynucleotide described in SEQ ID NO: 9.
2. An isolated nucleic acid molecule comprising a nucleotide sequence that:
 - (a) encodes the amino acid sequence shown in SEQ ID NO: 10; and
 - (b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO: 9 or the complement thereof.
3. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence disclosed in SEQ ID NO: 12.

SEQUENCE LISTING

<110> Donoho, Gregory
 Turner, C. Alexander Jr.
 Nehls, Michael
 Friedrich, Glenn
 Zambrowicz, Brian
 Sands, Arthur T.

<120> Novel Human Carboxypeptidases and
 Polynucleotides Encoding the Same

<130> LEX-0047-PCT

<150> US 60/156,685

<151> 1999-09-29

<160> 13

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 141

<212> DNA

<213> homo sapiens

<400> 1

atgaagtgtc	tcgggaagcg	caggggccag	gcagctgctt	tctgacctct	ttgctggctc	60
tttttgaaga	ttctgcaacc	ggggcacagc	cacctttata	acaaccgcta	tgctgggtcca	120
cagggaaaga	aacttttgaa	a				141

<210> 2

<211> 47

<212> PRT

<213> homo sapiens

<400> 2

Met	Lys	Cys	Leu	Gly	Lys	Arg	Arg	Gly	Gln	Ala	Ala	Ala	Phe	Leu	Pro
1			5					10					15		
Leu	Cys	Trp	Leu	Phe	Leu	Lys	Ile	Leu	Gln	Pro	Gly	His	Ser	His	Leu
			20				25					30			
Tyr	Asn	Asn	Arg	Tyr	Ala	Gly	Pro	Gln	Gly	Lys	Lys	Leu	Leu	Lys	
	35					40					45				

<210> 3

<211> 264

<212> DNA

<213> homo sapiens

<400> 3

atgaagtgtc	tcgggaagcg	caggggccag	gcagctgctt	tctgacctct	ttgctggctc	60
tttttgaaga	ttctgcaacc	ggggcacagc	cacctttata	acaaccgcta	tgctgggtgat	120
aaagtgataa	gattttattcc	caaaacagaa	gaggaagcat	atgcactgaa	gaaaatatcc	180
tatcaactta	aggttggttc	ctgcactaca	ggtggacctg	tggcagccca	gcagtatctc	240

ctatgtatca gaggggaacag ttac

264

<210> 4

<211> 88

<212> PRT

<213> homo sapiens

<400> 4

Met Lys Cys Leu Gly Lys Arg Arg Gly Gln Ala Ala Ala Phe Leu Pro
 1 5 10 15
 Leu Cys Trp Leu Phe Leu Lys Ile Leu Gln Pro Gly His Ser His Leu
 20 25 30
 Tyr Asn Asn Arg Tyr Ala Gly Asp Lys Val Ile Arg Phe Ile Pro Lys
 35 40 45
 Thr Glu Glu Glu Ala Tyr Ala Leu Lys Lys Ile Ser Tyr Gln Leu Lys
 50 55 60
 Val Gly Ser Cys Thr Thr Gly Gly Pro Val Ala Ala Gln Gln Tyr Leu
 65 70 75 80
 Leu Cys Ile Arg Gly Asn Ser Tyr
 85

<210> 5

<211> 741

<212> DNA

<213> homo sapiens

<400> 5

atgaagtgtc tcgggaagcg cagggggccag gcagctgctt tcctgcctct ttgctggctc 60
 tttttgaaga ttctgcaacc ggggcacagc cacctttata acaaccgcta tgctgggtgat 120
 aaagtgataa gatttattcc caaaacagaa gaggaagcat atgcactgaa gaaaatatcc 180
 tatcaactta aggtggacct gtggcagccc agcagtatct cctatgtatc agaggggaaca 240
 gttactgatg tccatatccc ccaaaatggt tcccagagccc tgttagcctt cttacaggaa 300
 gccaacatcc agtacaaggt cctcatagaa gatcttcaga aaacactgga gaaggggaagc 360
 agcttgacaca cccagagaaa ccgaagatcc ctctctggat ataattatga agtttatcac 420
 tccttagaag aaattcaaaa ttggatgcat catctgaata aaactcactc aggcctcatt 480
 cacatgttct ctattggaag atcatatgag ggaagatctc tttttatttt aaagctgggc 540
 agacgatcac gactcaaaaag agctgtttgg atagactgtg gtattcatgc aagagaatgg 600
 attggctcctg ccttttgtca gtggtttgta aaagaagtc tagaaaaacac agctcacaaa 660
 tgtcaagaat gtactaaatt taaaaaatat ctctgccact accaaaacca caaaagtatg 720
 cttaatcttg taagtattga g 741

<210> 6

<211> 247

<212> PRT

<213> homo sapiens

<400> 6

Met Lys Cys Leu Gly Lys Arg Arg Gly Gln Ala Ala Ala Phe Leu Pro
 1 5 10 15
 Leu Cys Trp Leu Phe Leu Lys Ile Leu Gln Pro Gly His Ser His Leu
 20 25 30
 Tyr Asn Asn Arg Tyr Ala Gly Asp Lys Val Ile Arg Phe Ile Pro Lys
 35 40 45
 Thr Glu Glu Glu Ala Tyr Ala Leu Lys Lys Ile Ser Tyr Gln Leu Lys
 50 55 60
 Val Asp Leu Trp Gln Pro Ser Ser Ile Ser Tyr Val Ser Glu Gly Thr

65		70		75		80
Val Thr Asp Val	His Ile Pro Gln Asn Gly Ser Arg Ala Leu Leu Ala					
	85		90			95
Phe Leu Gln Glu Ala Asn Ile Gln Tyr Lys Val Leu Ile Glu Asp Leu						
	100		105			110
Gln Lys Thr Leu Glu Lys Gly Ser Ser Leu His Thr Gln Arg Asn Arg						
	115		120			125
Arg Ser Leu Ser Gly Tyr Asn Tyr Glu Val Tyr His Ser Leu Glu Glu						
	130		135			140
Ile Gln Asn Trp Met His His Leu Asn Lys Thr His Ser Gly Leu Ile						
145		150		155		160
His Met Phe Ser Ile Gly Arg Ser Tyr Glu Gly Arg Ser Leu Phe Ile						
	165		170			175
Leu Lys Leu Gly Arg Arg Ser Arg Leu Lys Arg Ala Val Trp Ile Asp						
	180		185			190
Cys Gly Ile His Ala Arg Glu Trp Ile Gly Pro Ala Phe Cys Gln Trp						
	195		200			205
Phe Val Lys Glu Val Leu Glu Asn Thr Ala His Lys Cys Gln Glu Cys						
	210		215			220
Thr Lys Phe Thr Lys Tyr Leu Cys His Tyr Gln Asn His Lys Ser Met						
225		230		235		240
Leu Asn Leu Val Ser Ile Glu						
	245					

<210> 7

<211> 276

<212> DNA

<213> homo sapiens

<400> 7

atgaagtgtc	tcgggaagcg	cagggggccag	gcagctgctt	tcttgcctct	ttgctggctc	60
tttttgaaga	ttctgcaacc	ggggcacagc	cacctttata	acaaccgcta	tgctgggtgat	120
aaagtgataa	gattttattcc	caaaacagaa	gaggaagcat	atgcactgaa	gaaaatatcc	180
tatcaactta	agggtcctca	tagaagatct	tcagaaaaca	ctggagaagg	gaagcagctt	240
gcacacccag	agaaaccgaa	gatccctctc	tgata			276

<210> 8

<211> 92

<212> PRT

<213> homo sapiens

<400> 8

Met Lys Cys Leu Gly Lys Arg Arg Gly Gln Ala Ala Ala Phe Leu Pro		
1	5	10
Leu Cys Trp Leu Phe Leu Lys Ile Leu Gln Pro Gly His Ser His Leu		
	20	25
Tyr Asn Asn Arg Tyr Ala Gly Asp Lys Val Ile Arg Phe Ile Pro Lys		
	35	40
Thr Glu Glu Glu Ala Tyr Ala Leu Lys Lys Ile Ser Tyr Gln Leu Lys		
	50	55
Gly Pro His Arg Arg Ser Ser Glu Asn Thr Gly Glu Gly Lys Gln Leu		
65	70	75
Ala His Pro Glu Lys Pro Lys Ile Pro Leu Trp Ile		
	85	90

<210> 9

<211> 1311

<212> DNA

<213> homo sapiens

<400> 9

```

atgaagtgtc tcggaagcg cagggggccag gcagctgctt tcctgcctct ttgctggctc      60
tttttgaaga ttctgcaacc ggggcacagc cacctttata acaaccgcta tgctgggtgat      120
aaagtgataa gatttattcc caaaacagaa gaggaagcat atgcactgaa gaaaatatcc      180
tatcaactta aggtggacct gtggcagccc agcagtatct cctatgtatc agaggggaaca      240
gttactgatg tccatatccc ccaaaatggt tcccgagccc tgtagcctt cttacaggaa      300
gccaacatcc agtacaaggt cctcatagaa gatcttcaga aaactctgga gaagggaagc      360
agcttgacac cccagagaaa ccgaagatcc ctctctggat ataattatga agtttatcac      420
tccttagaag aaattcaaaa ttggatgcat catctgaata aaactcactc aggcctcatt      480
cacatgttct ctattggaag atcatatgag ggaagatctc tttttatttt aaagctgggc      540
agacgatcac gactcaaaag agctggttgg atagactgtg gtattcatgc aagagaatgg      600
attggtcctg ccttttgtca gtggtttgta aaagaagctc ttctaacata taagagtgc      660
ccagccatga gaaaaatggt gaatcatcta tatttctata tcatgcctgt gtttaacgtc      720
gatggatacc attttagtgt gaccaatgat cgattttgga gaaaaacaag gtcaaggaa      780
tcaaggtttc gctgccgtgg agtggatgcc aatagaaact ggaaagtga gtggtgtgat      840
gaaggagctt ctatgcaccc ttgtgatgac acatactgtg gcccttttcc agaactctgag      900
ccggaagtga aggtctgtagc taacttctt cgaaaacaca gaaagcacat tagggcttat      960
ctctcctttc atgcatatgc tcagatgtta ctgtatccct attcttacia atatgcaaca     1020
attcccaatt ttagatgtgt ggaatctgca gcttataaag ctgtgaatgc acttcagtca     1080
gtatacgggg tacgatacag atatggacca gcctocacaa cgttgtatgt gagctctggt     1140
agctcaatgg attgggccta caaaaatgga ataccttatg catttgcttt cgaactacgt     1200
gacactggat attttggtt tttactccca gagatgctca tcaaaccac ctgtacagaa     1260
actatgctgg ctgtgaaaaa tatcacaatg cacctgctaa agaatgtcc c              1311

```

<210> 10

<211> 437

<212> PRT

<213> homo sapiens

<400> 10

```

Met Lys Cys Leu Gly Lys Arg Arg Gly Gln Ala Ala Ala Phe Leu Pro
 1          5          10          15
Leu Cys Trp Leu Phe Leu Lys Ile Leu Gln Pro Gly His Ser His Leu
 20          25          30
Tyr Asn Asn Arg Tyr Ala Gly Asp Lys Val Ile Arg Phe Ile Pro Lys
 35          40          45
Thr Glu Glu Glu Ala Tyr Ala Leu Lys Lys Ile Ser Tyr Gln Leu Lys
 50          55          60
Val Asp Leu Trp Gln Pro Ser Ser Ile Ser Tyr Val Ser Glu Gly Thr
 65          70          75          80
Val Thr Asp Val His Ile Pro Gln Asn Gly Ser Arg Ala Leu Leu Ala
 85          90          95
Phe Leu Gln Glu Ala Asn Ile Gln Tyr Lys Val Leu Ile Glu Asp Leu
100          105          110
Gln Lys Thr Leu Glu Lys Gly Ser Ser Leu His Thr Gln Arg Asn Arg
115          120          125
Arg Ser Leu Ser Gly Tyr Asn Tyr Glu Val Tyr His Ser Leu Glu Glu
130          135          140
Ile Gln Asn Trp Met His His Leu Asn Lys Thr His Ser Gly Leu Ile
145          150          155          160
His Met Phe Ser Ile Gly Arg Ser Tyr Glu Gly Arg Ser Leu Phe Ile
165          170          175

```

Leu Lys Leu Gly Arg Arg Ser Arg Leu Lys Arg Ala Val Trp Ile Asp
 180 185 190
 Cys Gly Ile His Ala Arg Glu Trp Ile Gly Pro Ala Phe Cys Gln Trp
 195 200 205
 Phe Val Lys Glu Ala Leu Leu Thr Tyr Lys Ser Asp Pro Ala Met Arg
 210 215 220
 Lys Met Leu Asn His Leu Tyr Phe Tyr Ile Met Pro Val Phe Asn Val
 225 230 235 240
 Asp Gly Tyr His Phe Ser Trp Thr Asn Asp Arg Phe Trp Arg Lys Thr
 245 250 255
 Arg Ser Arg Asn Ser Arg Phe Arg Cys Arg Gly Val Asp Ala Asn Arg
 260 265 270
 Asn Trp Lys Val Lys Trp Cys Asp Glu Gly Ala Ser Met His Pro Cys
 275 280 285
 Asp Asp Thr Tyr Cys Gly Pro Phe Pro Glu Ser Glu Pro Glu Val Lys
 290 295 300
 Ala Val Ala Asn Phe Leu Arg Lys His Arg Lys His Ile Arg Ala Tyr
 305 310 315 320
 Leu Ser Phe His Ala Tyr Ala Gln Met Leu Leu Tyr Pro Tyr Ser Tyr
 325 330 335
 Lys Tyr Ala Thr Ile Pro Asn Phe Arg Cys Val Glu Ser Ala Ala Tyr
 340 345 350
 Lys Ala Val Asn Ala Leu Gln Ser Val Tyr Gly Val Arg Tyr Arg Tyr
 355 360 365
 Gly Pro Ala Ser Thr Thr Leu Tyr Val Ser Ser Gly Ser Ser Met Asp
 370 375 380
 Trp Ala Tyr Lys Asn Gly Ile Pro Tyr Ala Phe Ala Phe Glu Leu Arg
 385 390 395 400
 Asp Thr Gly Tyr Phe Gly Phe Leu Leu Pro Glu Met Leu Ile Lys Pro
 405 410 415
 Thr Cys Thr Glu Thr Met Leu Ala Val Lys Asn Ile Thr Met His Leu
 420 425 430
 Leu Lys Lys Cys Pro
 435

<210> 11
 <211> 1050
 <212> DNA
 <213> homo sapiens

<400> 11
 atgaagtgtc tcgggaagcg caggggccag gcagctgctt tcctgcctct ttgctggctc 60
 tttttgaaga ttctgcaacc ggggcacagc cacctttata acaaccgcta tgctggtgat 120
 aaagtgataa gatttattcc caaaacagaa gaggaagcat atgcactgaa gaaaatatcc 180
 tatcaactta aggtggaccc gtggcagccc agcagtatct cctatgtatc agagggaaca 240
 gttactgatg tccatatccc ccaaaatggt tcccagagccc tgttagcctt cttacaggaa 300
 gccaacatcc agtacaaggt cctcatagaa gatcttcaga aaacactgga gaagggaagc 360
 agcttgacaca cccagagaaa ccgaagatcc ctctctggat ataattatga agtttatcac 420
 tccttagaag aaattcaaaa ttggatgcat catctgaata aaactcactc aggcctcatt 480
 cacatgttct ctattggaag atcatatgag ggaagatctc tttttatttt aaagctgggc 540
 agacgatcac gactcaaaag agctgtttgg atagactgtg gtattcatgc aagagaatgg 600
 attggtcctg ccttttgtca gtggtttgta aaagaagctc ttctaacata taagagtgc 660
 ccagccatga gaaaaatggt gaatcatcta tatttctata tcatgcctgt gtttaacgtc 720
 gatggatacc attttagttg gaccaatgat cgattttgga gaaaaacaag gtcaaggaac 780
 tcaaggtttc gctgccgtgg agtggatgcc aatagaaact ggaaagtgaa gtggtgtgat 840
 gaaggagctt ctatgcaccc ttgtgatgac acatactgtg gcccttttcc agaactctgag 900

ccggaagtga aggctgtagc taacttcctt cgaaaacaca gaaagcacat tagggcttat 960
 ctctcctttc atgcatatgc tcagatgtta ctgtatccct attcttacaa atatgcaaca 1020
 attccaatt ttagatgtgt ggtaagtatt 1050

<210> 12
 <211> 350
 <212> PRT
 <213> homo sapiens

<400> 12

Met Lys Cys Leu Gly Lys Arg Arg Gly Gln Ala Ala Ala Phe Leu Pro
 1 5 10 15
 Leu Cys Trp Leu Phe Leu Lys Ile Leu Gln Pro Gly His Ser His Leu
 20 25 30
 Tyr Asn Asn Arg Tyr Ala Gly Asp Lys Val Ile Arg Phe Ile Pro Lys
 35 40 45
 Thr Glu Glu Glu Ala Tyr Ala Leu Lys Lys Ile Ser Tyr Gln Leu Lys
 50 55 60
 Val Asp Leu Trp Gln Pro Ser Ser Ile Ser Tyr Val Ser Glu Gly Thr
 65 70 75 80
 Val Thr Asp Val His Ile Pro Gln Asn Gly Ser Arg Ala Leu Leu Ala
 85 90 95
 Phe Leu Gln Glu Ala Asn Ile Gln Tyr Lys Val Leu Ile Glu Asp Leu
 100 105 110
 Gln Lys Thr Leu Glu Lys Gly Ser Ser Leu His Thr Gln Arg Asn Arg
 115 120 125
 Arg Ser Leu Ser Gly Tyr Asn Tyr Glu Val Tyr His Ser Leu Glu Glu
 130 135 140
 Ile Gln Asn Trp Met His His Leu Asn Lys Thr His Ser Gly Leu Ile
 145 150 155 160
 His Met Phe Ser Ile Gly Arg Ser Tyr Glu Gly Arg Ser Leu Phe Ile
 165 170 175
 Leu Lys Leu Gly Arg Arg Ser Arg Leu Lys Arg Ala Val Trp Ile Asp
 180 185 190
 Cys Gly Ile His Ala Arg Glu Trp Ile Gly Pro Ala Phe Cys Gln Trp
 195 200 205
 Phe Val Lys Glu Ala Leu Leu Thr Tyr Lys Ser Asp Pro Ala Met Arg
 210 215 220
 Lys Met Leu Asn His Leu Tyr Phe Tyr Ile Met Pro Val Phe Asn Val
 225 230 235 240
 Asp Gly Tyr His Phe Ser Trp Thr Asn Asp Arg Phe Trp Arg Lys Thr
 245 250 255
 Arg Ser Arg Asn Ser Arg Phe Arg Cys Arg Gly Val Asp Ala Asn Arg
 260 265 270
 Asn Trp Lys Val Lys Trp Cys Asp Glu Gly Ala Ser Met His Pro Cys
 275 280 285
 Asp Asp Thr Tyr Cys Gly Pro Phe Pro Glu Ser Glu Pro Glu Val Lys
 290 295 300
 Ala Val Ala Asn Phe Leu Arg Lys His Arg Lys His Ile Arg Ala Tyr
 305 310 315 320
 Leu Ser Phe His Ala Tyr Ala Gln Met Leu Leu Tyr Pro Tyr Ser Tyr
 325 330 335
 Lys Tyr Ala Thr Ile Pro Asn Phe Arg Cys Val Val Ser Ile
 340 345 350

<210> 13

<211> 2128

<212> DNA

<213> homo sapiens

<400> 13

attaaagatc	aggtcagctg	ctgctgctgc	tgctgctgct	tgtcccaaga	ccaagtcgta	60
atagcaactt	cccttcctca	gctgcctgaa	cttttttttt	cccttgtagc	tggagagaag	120
tgtcacattt	tgctcactct	caaccttcct	cgccccaccc	cttcccggag	aacctgtgcg	180
gtgtgtagag	ggtgctgtga	gccacctcca	gcctcgggtg	gctgcttaag	taactttcaa	240
ctcctctctt	cttaacaacta	tgaagtgtct	cgggaagcgc	agggggccagg	cagctgcttt	300
cctgcctctt	tgctggctct	ttttgaagat	tctgcaaccg	gggcacagcc	acctttataa	360
caaccgctat	gctggtccac	agggaaagaa	acttttgaaa	taataatcag	ttgcctggta	420
tactgctcaa	tgatattgcc	acacgtaaga	acaagcatac	agtataaag	tgataagatt	480
tattcccaaa	acagaagagg	aagcatatgc	actgaagaaa	atatcctatc	aacttaagggt	540
tggttcctgc	actacaggtg	gacctgtggc	agcccagcag	tatctcctat	gtatcagagg	600
gaacagttac	tgatgtccat	atcccccaaa	atggttcccg	agccctgtta	gccttcttac	660
aggaagccaa	catccagtac	aaggctcctca	tagaagatct	tcagaaaaca	ctggagaagg	720
gaagcagctt	gcacacccag	agaaaccgaa	gatccctctc	tggatataat	tatgaagttt	780
atcactcctt	agaagaaatt	caaaattgga	tgcatcatct	gaataaaaact	cactcaggcc	840
tcattcacat	gttctctatt	ggaagatcat	atgagggaag	atctcttttt	attttaaagc	900
tgggcagacg	atcacgactc	aaaagagctg	tttgataga	ctgtggtatt	catgcaagag	960
aatggattgg	tcctgccttt	tgtcagtggt	ttgtaaaaga	agtcctagaa	aacacagctc	1020
acaaatgtca	agaatgtact	aaattttacaa	aatatctctg	ccactaccaaa	aaccacaaaa	1080
gtatgcttaa	tcttgtaagt	attgagtaat	aaaattttct	aaacattcct	aaaaactcct	1140
ctaacaata	agagtgacct	agccatgaga	aaaatggtga	atcatctata	tttctatata	1200
atgcctgtgt	ttaacgtcga	tggataccat	tttagttgga	ccaatgatcg	attttggaga	1260
aaaacaaggt	caaggaactc	aagggttcgc	tgccgtggag	tggatgccaa	tagaaactgg	1320
aaagtgaagt	ggtgtgatga	aggagcttct	atgcaccctt	gtgatgacac	atactgtggc	1380
ccttttccag	aatctgagcc	ggaagtgaag	gctgtagcta	acttccttcg	aaaacacaga	1440
aagcacatta	gggcttatct	ctcctttcat	gcatatgctc	agatgttact	gtatccctat	1500
tottacaaat	atgcaacaat	tcccaatttt	agatgtgtgg	aatctgcagc	ttataaagct	1560
gtgaatgcac	ttcagtcagt	atacggggta	cgatacagat	atggaccagc	ctccacaacg	1620
ttgtatgtga	gctctggtag	ctcaatggat	tgggcctaca	aaaatgggaat	accttatgca	1680
tttgctttcg	aactacgtga	cactggatat	tttggttttt	tactcccaga	gatgctcatc	1740
aaaccacact	gtacagaaac	tatgctggct	gtgaaaaata	tcacaatgca	cctgctaaag	1800
aaatgtccct	gagacagccc	aaggctcaag	tcaactgcc	taggattctg	agcaaggcct	1860
acttggccct	ggatagaaat	tgttttcaaa	gagaagggca	gctgcttaga	gtgaacatgt	1920
ctatggactt	taaaaagacc	ccacgcaatt	ttgacttttt	ttggggccaa	tttggaaaaa	1980
acagttaagt	atttgaccct	gtgcatgtac	atcaggcttc	atgctgcttt	tctgaagcta	2040
agatggttct	aagtactaat	gataatggca	aacacatggt	tgtgtttatc	ctaataaata	2100
ttttacatgt	gaaaaaaaaa	aaaaaaaaa				2128

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/26876

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/57 C12N9/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, STRAND

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL Database, Heidelberg, FRG Emest accession number H86718 22 November 1995 HILLIER, L. ET AL.: "yt08c10.r1 Soares retina N2b5HR Homo sapiens cDNA clone IMAGE: 223698 5', mRNA sequence" XP002162175	1
A	the whole document	2,3
A	WO 98 35988 A (ZENECA LIMITED) 20 August 1998 (1998-08-20) see especially SEQ ID NOS: 11 and 12 the whole document	1-3
A	WO 96 23064 A (BIOTECHNOLOGY GENERAL CORP.) 1 August 1996 (1996-08-01) see especially SEQ ID NOS: 5 and 6 the whole document	1-3
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

6 March 2001

Date of mailing of the international search report

19/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fuchs, U

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/26876

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	EMBL Database, Heidelberg, FRG Emhum2 accession number AF221594 2 August 2000 GU, J. ET AL.: "Homo sapiens carboxypeptidase B precursor (CPAH) mRNA, complete cds." XP002162176 the whole document	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/26876

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9835988 A	20-08-1998	AU 6000698 A	08-09-1998
WO 9623064 A	01-08-1996	AU 698889 B	12-11-1998
		AU 4903496 A	14-08-1996
		BR 9606795 A	30-12-1997
		CA 2210242 A	01-08-1996
		CN 1177377 A	25-03-1998
		CZ 9702258 A	14-10-1998
		EP 0871718 A	21-10-1998
		HU 9800091 A	28-05-1998
		IL 116696 A	17-08-1999
		JP 11503002 T	23-03-1999
		NZ 302874 A	25-11-1998
		PL 321499 A	08-12-1997
		US 5948668 A	07-09-1999
		ZA 9600515 A	15-08-1996